



SKIN & BEAUTY CENTER
Cosmetic, Medical and Surgical Dermatology

DermLA.com

BURBANK
191 S. Buena Vista St. Suite 475
Burbank, CA 91505
T: (818) 842-8000
F: (818) 842-3208

SANTA CLARITA
27879 Smyth Drive
Valencia, CA 91355
T: (877) 822-2223
F: (323) 935-8804



NORTHRIDGE
9535 Reseda Blvd. Suite 304
Northridge, CA 91324
T: (818) 886-3884
F: (818) 886-5418

William Coleman, MD
David Friedman, MD
Ava Khosraviani, MD
Payam Saadat, MD
Manjunath Vadmal, MD
Kristen Ochsner, PA-C
Jenna Trubschenck, M.S. PA-C

Erythema Nodosum

Erythema Nodosum is a form of 'panniculitis' or inflammation of subcutaneous fat tissue. It is quite painful, especially if pressed upon. It looks like painful bruises on both of the shins. A few other bumps may be present elsewhere, for example, the knees, elbows, forearms and thighs. You may also feel ill, like you have the flu. Erythema Nodosum is most likely to occur between fifteen and thirty years of age. Women are affected three times as often as men are.

They start out slightly raised and bright red. They can range in size from one half inch to several inches across. The color turns to a purplish red. As they start to fade they get a yellowish color.

There are many causes of Erythema Nodosum. The most common cause of Erythema Nodosum is medications, especially Sulfa drugs and some brands of the oral contraceptive pill. If the pill is changed, the Erythema Nodosum resolves. Apart from drugs, infections are the largest cause of Erythema Nodosum, including 'Strep' throat, TB and others.

The skin lesions of Erythema Nodosum are quite characteristic, nonetheless, they need to be differentiated from other conditions with similar symptoms. These include insect bites, bruising, pancreatitis and phlebitis. A biopsy is usually done, along with a blood test and a chest x-ray.

The most effective treatment is to remove the cause. Sometimes no cause is found, and complete resolution takes four to six weeks. Non-steroidal anti-inflammatory (NSAID's-Ibuprofen, Alleve and others) and bed rest may be used to reduce the pain and swelling. Injections may also be used for large and painful nodules. Oral potassium iodide works great but is unpleasant to take and must be dosed correctly. Other treatments are available in severe cases.